



Aquarius L3 Gridded 1-Degree Monthly Soil Moisture Climatology, Version 5

USER GUIDE

How to Cite These Data

As a condition of using these data, you must include a citation:

Bindlish, R. and T. Jackson. 2018. *Aquarius L3 Gridded 1-Degree Monthly Soil Moisture Climatology, Version 5*. [Indicate subset used]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. <https://doi.org/10.5067/O19KHW22C685>. [Date Accessed].

FOR QUESTIONS ABOUT THESE DATA, CONTACT NSIDC@NSIDC.ORG

FOR CURRENT INFORMATION, VISIT https://nsidc.org/data/AQ3_MCSM



National Snow and Ice Data Center

TABLE OF CONTENTS

1	DETAILED DATA DESCRIPTION.....	2
1.1	Format	2
1.2	File and Directory Structure.....	2
1.3	File Naming Convention	2
1.4	Volume	3
1.5	Spatial Coverage.....	3
1.5.1	Spatial Resolution	3
1.5.2	Projection and Grid Description	3
1.6	Temporal Coverage.....	3
1.6.1	Temporal Resolution.....	4
1.7	Parameter or Variable	4
1.7.1	Parameter Description	4
2	SOFTWARE AND TOOLS	5
3	DATA ACQUISITION AND PROCESSING.....	5
3.1	Theory of Measurements.....	5
3.2	Data Acquisition Methods.....	6
3.3	Derivation Techniques and Algorithms.....	6
3.3.1	Processing Steps	7
3.3.2	Version History.....	7
3.4	Sensor or Instrument Description.....	8
4	REFERENCES AND RELATED PUBLICATIONS	9
4.1	Related Data Collections.....	10
4.2	Related Websites	10
5	CONTACTS AND ACKNOWLEDGMENTS	10
6	DOCUMENT INFORMATION.....	11
6.1	Publication Date	11
6.2	Date Last Updated.....	11

1 DETAILED DATA DESCRIPTION

1.1 Format

Data are provided in Hierarchical Data Format 5 (HDF5) files. Data values are stored as bytes, 2-byte integers, and 4-byte floats. Soil moisture data are 32-bit float and color palette values are 8-bit unsigned integer. Each data file is paired with an associated XML file. XML files contain file level metadata and location, platform, and campaign information.

1.2 File and Directory Structure

Data are available at:

https://n5ei101u.ecs.nsidc.org/AQUARIUS/AQ3_MC.005/

Data files are organized in directories by date in YYYY.MM.DD format, for example:

/2011.08.25/

/2011.09.01/

/2011.10.01/

/2011.11.01/

1.3 File Naming Convention

All files are named according to the convention shown in the following example. File name variables are defined in Table 1.

Example

Q20112442014273.L3m_MC_SOILM_V5.0_rad_sm_1deg

Naming Convention

QYYYYDDDDyyyyddd.L3m_TT_SOILM_V5.0_rad_sm_1deg

Table 1. File Naming Convention

Variable	Description
Q	Indicates Aquarius instrument
YYYYDDD	Start date (4-digit year, 3-digit day)
yyyyddd	End date (4-digit year, 3-digit day)

Variable	Description
L3m	Processing level (Level-3 mapped)
TT	Observation period: MC: Monthly Climatology
SOILM	Parameter (SOILM: soil moisture)
V5.0	Data version number (V5.0)
rad_sm_1deg	1-degree radiometer soil moisture

1.4 Volume

The total volume for all Aquarius L3 monthly soil moisture climatology files is approximately 4 MB.

1.5 Spatial Coverage

The spatial coverage for this data set is global.

1.5.1 Spatial Resolution

Data are 1-degree spatial resolution.

1.5.2 Projection and Grid Description

The main HDF5 data group, called `l3m_data`, is a two-dimensional array (180 rows, 360 columns) of an Equidistant Cylindrical projection (also known as Plate Carrée) of the globe.

1.6 Temporal Coverage

The temporal coverage for this data set spans from 25 August 2011 through 31 May 2015.

Monthly climatology products contain the mean soil moisture values of composites for the Aquarius mission, including:

- 25 August 2011 to 31 August 2014
- 01 September 2011 to 30 September 2014
- 01 October 2011 to 31 October 2014
- 01 November 2011 to 30 November 2014
- 01 December 2011 to 31 December 2014
- 01 January 2012 to 31 January 2015
- 01 February 2012 to 28 February 2015
- 01 March 2012 to 31 March 2015
- 01 April 2012 to 30 April 2015
- 01 May 2012 to 31 May 2015

- 01 June 2012 to 30 June 2014
- 01 July 2012 to 31 July 2014

 Due to a power failure on the Satélite de Aplicaciones Científicas (SAC)-D spacecraft on 08 June 2015, data from the NASA Aquarius instrument were no longer being produced after 07 June 2015. Accordingly, the date range for the climatology Spring season ends on 07 June 2015, rather than extending through the full Spring season.

1.6.1 Temporal Resolution

The temporal resolution for this data set is monthly.

1.7 Parameter or Variable

This Level-3 product is a representation of a binned data product generated from Aquarius data. The HDF5 data group called 13m_data represents mean soil moisture at each grid point.

1.7.1 Parameter Description

Each Level-3 soil moisture file contains the group 13m_data with the attributes described in the following table.

Table 2. Attributes of the 13m_data Group

Name	Value
Scaling	linear
Scaling Equation ¹	(Slope*13m_data) + Intercept = Parameter value
Slope	1.0
Intercept	0
_FillValue	-32767.0
add_offset	0.0
scale_factor	1.0
¹ Scaling equations are typically provided to rescale the data for plotting purposes; however, in this case the scaling equation results in no change to the data values, and therefore can be disregarded.	

The group called palette in each data file is the color palette used in the graphics generated from the mapped files. This, or any other palette, can be used with these data.

Metadata are included as global attributes within each data file and have been adapted for Version 5 to more closely align with Climate and Forecast (CF) metadata conventions. A total of 68

metadata fields are provided, such as minimum and maximum data values, units, and platform and projection information. Values for some data fields may vary from granule to granule.

Figure 1 shows the monthly soil moisture climatology estimates for April 2015.




Figure 1. This plot shows Aquarius monthly soil moisture climatology estimates acquired using all three beams for April 2015.

2 SOFTWARE AND TOOLS

The following resources provide access to software for reading and viewing HDF5 data files:

- [HDFView](#) — Visual tool for browsing and editing HDF4 and HDF5 files; additional tools are available on the [HDF5 Tools and Software website](#).
- [Panoply, NetCDF, HDF, and GRIB Data Viewer](#)— Cross-platform application that plots geo-gridded arrays from NetCDF, HDF, and GRIB data sets.

3 DATA ACQUISITION AND PROCESSING

3.1 Theory of Measurements

The Aquarius Single Channel Algorithm (SCA) uses the L-band horizontally polarized (h-pol) brightness temperature observations due to the higher sensitivity of this channel to soil moisture.

The Aquarius SCA approach is based on the simplified radiative transfer model developed under the assumption that the canopy and soil temperatures are the same (Jackson 1993). The SCA is applied to the individual Aquarius footprint Level-2 brightness temperature observations to produce a swath-based time-ordered product. (Bindlish and Jackson 2013; Bindlish et al. 2013).

3.2 Data Acquisition Methods

This Version 5 Aquarius L3 soil moisture data set is generated from brightness temperature measurements included in the [NASA Aquarius Level-2 Sea Surface Salinity & Wind Speed Data V5.0](#) product. The best quality data are selected for each orbit during Level-0 (L0) to Level-1A (L1A) data processing and are then used to create the L2 file that is input to the L3 science file.

3.3 Derivation Techniques and Algorithms

The Aquarius Level-3 gridding algorithm uses local polynomial fitting to grid the Level-2 soil moisture retrievals on a 1 degree grid (Fan and Gijbels, 1996; Lilly and Lagerloef, 2008). The Level-3 processing of Aquarius satellite data takes measurements at the boresight locations of the three radiometer beams, which have been already converted into physical units of soil moisture, and maps these onto a 1 degree grid.

This method fits a Pth-order polynomial at each grid point x_m . For data values g_n observed at locations x_n , $n = 1, 2, \dots N$, this corresponds to minimizing

$$\sum_{n=1}^N \left| g_n - \sum_{p=0}^P \widehat{\beta}_p(x) [x_n - x]^2 \right|^2 K_h(x_n - x) \tag{Equation 1}$$

at every grid point $x = x_m$, where

$$K_h(x) = K\left(\frac{x}{h}\right) / h \tag{Equation 2}$$

is a decaying weighting function which depends upon the bandwidth h , with $K(x)$ being a probability distribution function.

The regression coefficients

$$\widehat{\beta}_p(x) \tag{Equation 3}$$

$P = 1, 2, \dots P$ vary with spatial location, and are estimated at all grid point locations.

The function $g(x)$ is estimated by the lowest order coefficient,

$$\hat{g}(x) = \hat{\beta}_0(x) \quad \text{(Equation 4)}$$

while higher-order regression coefficients estimate the derivatives of the field through

$$g^{(p)}(x) = p! \hat{\beta}_{p0}(x) \quad \text{(Equation 5)}$$

The above discussion focuses on a 1-dimensional application, but can be extended to a 2-dimensional application. A complete description for the 2-dimensional problem is available in Fan and Gijbels (1996) and Lilly and Lagerloef (2008).

3.3.1 Processing Steps

Aquarius L3 Gridded 1-Degree Soil Moisture Climatology data are produced by NASA Goddard Space Flight Center's Ocean Data Processing System (ODPS).

This product represents data binned over the period covered by the input Aquarius product. The mean for the observation period is used to obtain the values for the grid points from the binned data products. Each product contains one soil moisture image and is stored in one physical HDF5 file. The data are not filtered during the gridding process. The user is advised to refer to the flags in the [Aquarius L2 Swath Single Orbit Soil Moisture Data](#).

Monthly climatology products are the mean value of all monthly composites for the mission, for example:

$$(\text{January 2012} + \text{January 2013} + \text{January 2014}) / 3$$

3.3.2 Version History

The following table outlines the version history for this product.

Table 3. Version History Details

Version	Description
V5	Changes to this version include: Updated Version 5 Aquarius brightness temperature data were used as input File-level metadata were modified to more closely align with Climate and Forecast (CF) metadata conventions

Version	Description
V4	For Version 4 Aquarius L3 Gridded 1-Degree Soil Moisture Data, the data set was updated to use the most recent version (Version 4) of Aquarius Brightness Temperatures as input.
V3	The Aquarius L3 Gridded 1-Degree Soil Moisture Data, Version 3 were processed from the Aquarius L2 Swath Single Orbit Soil Moisture Data, Version 3. Changes to this version include: Use of the most recent version (Version 3) of Aquarius Brightness Temperatures as input Aquarius Brightness Temperatures are no longer re-calibrated before soil moisture retrievals as was done for Version 2 data Soil moisture observations are valid over a wider range of brightness temperatures compared to Version 2 data Updates were made to the soil moisture model parameters (b and ω)
V2	First public data release

3.4 Sensor or Instrument Description

Aquarius/SAC-D is a collaboration between NASA and Argentina's space agency, Comisión Nacional de Actividades Espaciales (CONAE), with participation from Brazil, Canada, France and Italy. The Aquarius instrument was built jointly by NASA's Jet Propulsion Laboratory and NASA's Goddard Space Flight Center.

The Aquarius instrument includes three radiometers and one scatterometer. The soil moisture data are collected by the radiometers. The radiometers measure brightness temperature at 1.414 GHz in the horizontal and vertical polarizations (T_H and T_V). The scatterometer is a microwave radar sensor that measures backscatter for surface roughness corrections. Table 4 summarizes instrument characteristics.

Table 4. Aquarius Instrument Characteristics

Instrument	Characteristics
3 radiometers in push-broom alignment	<ul style="list-style-type: none"> • Frequency: 1.413 GHz • Band width: less than or equal to 26 MHz • Swath Width: 390 km • Science data block period: 1.44 sec • Footprints for the beams are: 74 km along track x 94 km cross track, 84 x 120 km, and 96 x 156 km, yielding a total cross track of 390 km. • Beam incidence angles of 29.36, 38.49, and 46.29 degrees incident to the surface. Beams point away from the sun.

Instrument	Characteristics
Scatterometer	<ul style="list-style-type: none"> • Frequency: 1.26 GHz • Band Width: 4 MHz • Swath Width: 390 km • Science data block period: 1.44 sec

SAC-D spacecraft Orbit Parameters:

- 98 minute sun-synchronous
- 6 PM ascending orbit, 6 AM descending orbit
- 657 km equatorial altitude (655 km minimum, 685 km maximum over the orbit)
- Ground-track repeat interval: Weekly, 103 orbits

4 REFERENCES AND RELATED PUBLICATIONS

Bindlish, Rajat, and Thomas J. Jackson. 2013. Aquarius Soil Moisture ATBD Users Guide, Version 2.0. Beltsville, Maryland USA: USDA Hydrology and Remote Sensing Lab.

(<https://nsidc.org/sites/nsidc.org/files/files/data/aquarius/Aquarius-VSM-ATBD-UsersGuide.pdf>, 315 KB)

Bindlish, Rajat, Thomas Jackson, Michael Cosh, Tianjie Zhao and Peggy O'Neill. 2015. Global Soil Moisture from the Aquarius Satellite: Description and Initial Assessment. *IEEE Geosciences and Remote Sensing Letters* 12(5):923-927.

Fan, J. and I. Gijbels. 1996. *Local Polynomial Modelling and its Applications*, Chapman and Hall, 1996.

Jackson, T. J. 1993. Measuring Surface Soil Moisture Using Passive Microwave Remote Sensing. *Hydrological Processes* 7:139–152.

Lilly, Jonathan and Gary Lagerloef. 2008. *Aquarius Level 3 Processing Algorithm Theoretical Basis Document*. ftp://podaac-ftp.jpl.nasa.gov/allData/aquarius/docs/v2/AquariusLevel3_GriddingSmoothingPaper_Lilly&Lagerloef2008.pdf

Piepmeyer, Jeffrey, Shannon Brown, Joel Gales, Liang Hong, Gary Lagerloef, David Le Vine, Paolo de Mattheais, Thomas Meissner, Rajat Bindlish, and Thomas Jackson. 2013. *Aquarius Radiometer Post-Launch Calibration for Product Version 2.0*, Aquarius Project Document: AQ-014-PS-0015. ftp://podaac-ftp.jpl.nasa.gov/allData/aquarius/docs/v2/AQ-014-PS-0015_AquariusInstrumentCalibrationDescriptionDocument.pdf.

4.1 Related Data Collections

[AMSR-E/Aqua L2B Surface Soil Moisture, Ancillary Params, & QC EASE-Grids, Version 2](#)

[SMAP Data Sets at NSIDC](#)

[AMSR-E/Aqua Daily L3 Surface Soil Moisture, Interpretive Parameters, & QC EASE-Grids, Version 2](#)

[AMSR-E Validation Soil Moisture Data](#)

[Aquarius Level-1 and Level-2 Sea Surface Salinity Data](#)

[Aquarius Level-2 Swath Single Orbit Soil Moisture Data](#)

[ESA Soil Moisture and Ocean Salinity \(SMOS\)](#)

[Soil Moisture Product Using Aquarius/SAC-D Observations](#)

4.2 Related Websites

[Aquarius L2 Soil Moisture Documentation](#)

[Aquarius Web site at NASA Goddard Space Flight Center](#)

[Aquarius Data Web Site at NSIDC](#)

[Aquarius Web Site at PODAAC](#)

[SMAP Web Site at NSIDC](#)

[SMOS Website at ESA](#)

[NASA Aquarius Gallery: Soil Moisture - monthly soil moisture map images](#)

5 CONTACTS AND ACKNOWLEDGMENTS

Rajat Bindlish

NASA Goddard Space Flight Center

Hydrological Sciences Laboratory

Code 617, Bldg 33, G216

Greenbelt, MD 20771 USA

Thomas Jackson

United States Department of Agriculture

Agricultural Research Service

Hydrology and Remote Sensing Laboratory

Beltsville, MD 20705 USA

Acknowledgments:

This work was funded by NASA under the Interagency agreement NNH10AN10I. Tianjie Zhao helped with development of the soil moisture algorithm. The support provided by Michael Cosh,

Peggy O'Neill, Thomas Holmes and Wade Crow is acknowledged. We acknowledge the support provided by Gary Lagerloef, David Le Vine, Gene Feldman and the Aquarius Data Processing Segment (ADPS) group in the implementation of the Aquarius Soil moisture algorithm.

6 DOCUMENT INFORMATION

6.1 Publication Date

October 2015

6.2 Date Last Updated

September 2018